

# Micropropagation and Conservation of a Wild Species of *Solanum* through Organ Cultures

## Research Article

Salahuddin K<sup>1\*</sup>, Singh CK<sup>2</sup>, Md. Nizamuddin A<sup>2</sup> and Naseem M<sup>2</sup>

<sup>1</sup>Department of Botany, L.N. Mithila University, India

<sup>2</sup>University Department of Botany, B.R.A Bihar University, India

\*Corresponding author: Salahuddin K, Department of Botany, M.R.M College, L.N. Mithila University, Darbhanga-846004, Bihar, India, Phone: 9668816209, E-mail: salahuddin212@gmail.com

**Copyright:** © Salahuddin K, et al. 2020. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Article Information:** Submission: 10/03/2020; Accepted: 02/04/2020; Published: 10/04/2020

### Abstract

Micropropagation technique has emerged as the best tool for mass propagation of desired species. *Solanum torvum* Swartz, a wild species of crop plants is in high demand for medicinal as well as breeding point of view. The existence of these wild plants is in danger due to induction of new cultivars and other environmental hazards. In this background, there is urgent need for germplasm conservation and germplasm improvement besides mass propagation. Keeping these objectives into consideration, tissue culture studies of this plant were being undertaken to develop protocol for *in vitro* mass propagation and callogenesis. Regeneration of shoots and callus was obtained using sterilized segments of node (8-10 mm), internode (10-15 mm) and shoot-tip (8-10 mm) of *Solanum torvum* (about 2 years old). These explants were cultured on MS medium containing 0.8% agar, 3% sucrose and different combinations and concentrations of NAA/2,4-D and Kinetin (Kn) to obtain regenerates / plantlets and callus differentiation. Techniques were used for shoot regeneration directly from node and shoot-tip explants as well as from callus. Shoot regeneration was best achieved on 3mg/L-Kn in nodal and shoot-tip cultures. Callus mediated shoot regeneration was promising in the culture and was obtained on 2 mg/L<sup>-1</sup> NAA and 4 mg/L<sup>-1</sup>Kn on sub-culture. 2,4-D alone or 2,4-D + Kn resulted in callus differentiation from explants. Callus in general was white/greenish-white, compact, hydrated and crystalline in appearance. Callus was maintained for about 2 years on 1 mg/L<sup>-1</sup> NAA and 1 mg/L<sup>-1</sup> Kn on regular sub-culture after 25 days. Callus turned brown on higher concentration (10mg/L<sup>-1</sup>) of auxin and Kn on sub-culture. Rooting of microshoots (about 5 cm) was obtained on RM (½MS Salts) containing 1mg/L<sup>-1</sup> NAA and 2 mg/L<sup>-1</sup> IBA. Plantlets were successfully transferred to soil and they survived well in nature. Explants taken during December to May were most regenerative. Plantlets obtained through *in vitro* were morphologically identical to parent plants. Nodal explants were superior to other explants (internode, shoot-tip) with respect to shoot regeneration, where as internodal explants was superior for callogenesis.

**Keywords:** Callus; Explant; Germplasm; *In vitro*; Regeneration

## Introduction

The primitive cultivars and wild relatives of crop plants constitute a pool of genetic diversity which is invaluable for future breeding programme. However, the existence of these plants is in danger due to induction of new cultivars and other environmental hazards. Germplasm includes plant parts which are used for maintenance, conservation and propagation of any biotype; it also acts as genetic pool.

Tissue culture is a method of *in vitro* culture of cell, tissue and organ in a sterile culture medium [1]. This technique can be referred

to as “Botanical laser” and its numerous uses are yet to be explored and fully understood. The tools of plant tissue culture are being applied to a wide range of biotechnology ventures and in particular to the clonal propagation and genetic up gradation of crop and medicinal plants [2]. In recent years, tissue culture techniques have become useful tools in the hands of plant scientists of all disciplines because these techniques are more handy, less time consuming and less labour involving over the conventional methods of breeding and propagation.

*Solanum torvum*, (Fam: Solanaceae) commonly known as Devil's fig is a bushy perennial wild plant measuring 150-300 cm in height

and usually growing in tropical and subtropical areas throughout the world as a weed of disturbed areas. It is found growing in pastures, road sides and wastelands but not significantly in cultivated land. It prefers moist and fertile soil and also tolerates drought and saline soils.

Fruits are eaten as vegetable and used as ingredient of pickles, it is said to be good for enlargement of the spleen. Fruits contain a number of potentially pharmacologically active chemicals including sapogenin, steroid, sterolin, chlorogenin and solasonine. The aqueous extracts of turkey berry (*Solanum. torvum*) were found lethal to mice or depressed the erythrocytes, leukocytes and platelets in their blood. Extracts of the plant are reported to be useful in the treatment of hyperactivity, colds and cough, pimples, skin diseases and leprosy [3]. This plant is also used medicinally for the treatment of epilepsy [4].

Conservation of germplasm of this wild crop is highly needed for developing perennial brinjal variety, a common vegetable for millions of people of the world and its medicinal uses are also required to be investigated in right perspectives. In this background, it is necessary to multiply this plant through *in vitro* methods. Calli and regenerates obtained through *in vitro* methods can be used for germplasm conservation as well as for biochemical analysis [5]. For rapid multiplication of these wild plants, micropropagation is being increasingly applied to supplement conventional methods of propagation.

Our investigation is based using explants collected from mature *in vivo* grown plant (about 2 years old) and the cultures were maintained under continuous, cool and white fluorescent light (2000 lux) during the whole experiment. In my opinion, the present investigation would be the first thorough studies on organ cultures of this taxon. As the tissues of mature plant are as a rule recalcitrant, the tissue culture studies with explants taken from mature plant are of great significance. Hence, the present studies were aimed at *in vitro* regeneration of *Solanum torvum* through direct and callus mediated shoot regeneration using explants taken from *in vivo* grown plant (about 2 years old) under different hormonal regimes. Attempts were also made to suggest methods for germplasm conservation through tissue culture.

## Materials and Methods

The experimental plant, *Solanum torvum* was procured from warm moist fertile areas and was subjected to tissue culture experiments with a view to exploring the possibilities of micropropagation protocol and genetic upgradation through the use of somaclonal variants among regenerated plants. This study was also aimed to develop protocol for *in vitro* conservation of germplasm. The methodology of tissue culture experiments includes the following steps:

- (i) Preparation of culture media
- (ii) Preparation of Explants
- (iii) Inoculation and Transfer
- (iv) Maintenance of Cultures
- (v) Effect of Seasonal Variation on Regeneration and
- (vi) Rooting and Transfer of Plantlets

Nutritional requirements for optimal growth of a tissue *in vitro* may vary from species to species. Even tissues from different parts of the same plant may have some specific requirements for their satisfactory growth [6]. A wide range of culture media differing in their elemental composition has been described in the literature. In the present study, MS medium were used as basal medium as this was suitable for regeneration and callus induction [7]. The medium was prepared as such:

- (i) Required quantities of agar (0.8% w/v) and sucrose (3% w/v) were weighed out.
- (ii) Sucrose was dissolved in some amount of distilled water to give a concentrated solution and was filtered through the Whatman filter paper No. 1 (9.0 cm) to remove the particulate impurities, if any
- (iii) Appropriate quantities of various stock solutions and growth regulators were added.
- (iv) Agar was dissolved in distilled water (in about ¼ of the final volume of the medium) by heating in a water bath. The dissolved agar solution & sucrose solution were mixed with stock solution.
- (v) The final volume of the medium was made upto 1 litre / required volume with distilled water.
- (vi) After proper mixing, the pH of the medium was adjusted to 5.8 using 0.1N NaOH or 0.1N HCl with the help of "Systronic" digital pH meter model no. 335.
- (vii) About 20 ml of the medium was poured into the culture tube (25 x 100 mm)
- (viii) The culture tubes were plugged with non-absorbent cotton wrapped in cheese cloth. The cotton plugs were wrapped with aluminium foils to prevent wetting during autoclaving.
- (ix) The culture vessels were transferred to appropriate baskets and autoclaved at 121°C for 20 minutes.
- (x) Slants were prepared by keeping the tubes tilted during cooling.

The pH of the medium was adjusted to 5.8 before autoclaving at 121°C for 20 minute. Various growth regulators and adjuvants used as supplement of the basal medium were IBA, NAA, 2,4-D & Kn. Stem segments (nodes and internodes) and leaf segments from youngest shoots and shoot-tip segments collected from *in vivo* grown mature plant (about 2 years old) of *Solanum torvum* during March to November were used as explants and were surface sterilized. These adjuvants were used in a wide range of concentration (1-10 mg l<sup>-1</sup>) either alone or in various combinations [8,9]. The stocks of various growth regulators were prepared. All the precautions were taken while sterilizing the tissues avoiding any damage to them. The following steps were undertaken for sterilization of tissues or organ explants.

- (i) Washing the explants in running tap water
- (ii) The explants were treated for 2 min. in 1% cetavlon

(cetrimide I.P. 20% w/v isopropyl alcohol B.P. 10% v/v) solution followed by thorough washing in running tap water.

- (iii) Washing and disinfecting the explants in 0.2%  $\text{HgCl}_2$  solution for 3 to 5 min. depending upon the nature of the explants.
- (iv) Further, washing them three or four times thoroughly with sterile distilled water in an aseptic condition and
- (v) Finally using sterile forcep, tissue explants were transferred to sterile Petri dishes and were cut into required size with sterile scalpel or blade. Usually, node and internode of 8-10 mm, leaf segments of 5x5 mm and shoot-tip 10-15 mm were trimmed out for explant preparation.

The cultures were incubated in culture room maintained at  $25 \pm 2^\circ\text{C}$  with a relative humidity of about 60% under continuous fluorescent light (2000 lux, cool and white).

Calli obtained from different explants were taken out of the culture tubes aseptically and kept in a presterilized. The callus was cut into several pieces of almost equal sizes with the help of a sterilized blade. Pieces of calli from the growing portions were inoculated into the culture tubes containing MS medium with different combinations and concentrations of growth regulators. The calli were incubated at  $25 \pm 2^\circ\text{C}$  for further growth and differentiation.

Microshoots (3-4 cm) obtained from shoot-tip, nodal segment and regenerative callus in *Solanum. torvum* were cultured on MS and rooting media (1/2 MS salts + full strength vitamins & amino acid) supplemented with IBA and NAA singly and in combination for rhizogenesis. Culture conditions were kept constant as in shoot regeneration (Temp.  $25 \pm 2^\circ\text{C}$ , Light - 2000 lux, continuous, cool, white and fluorescent).

## Results and Discussion

In the present experimental system, nodal segment (8-10 mm), internodal segment (8-10mm), shoot-tip (10-15 mm) and leaf (5x5 mm) of these sizes were taken for experimentation and these explants yielded better results in culture. It was also remarkable in the present system that a proper amount of growing callus was essential for inoculum to had better differentiation and regeneration, a small piece of callus having few cells could not survive in culture.

The composition of culture medium is the most important factor for the establishment of tissue culture. It is confirmed that there is no fixed combination of the medium which is suitable for all the plants and even the different organs of the same plant. A particular combination of the nutrient medium is suitable for a certain group of plants but the same combination proves ineffective for other plants [10,11]. So, the selection of proper culture medium is essential for the tissue culture experiment of any plant. The response of two basal media viz. MS and Nitsch was tested in case of present experimental system and the results have been presented. MS medium was found most suitable for shoot regeneration and callus growth. MS medium was proved equally well in many other plants too. Normally, a high cytokinin to auxin ratio promotes shoot formation while a higher auxin to cytokinin ratio favours root differentiation [12]. In a number of cases, cytokinin alone is sufficient for shoot regeneration and callus

formation [13]. Identical response of cytokinin was encountered in *Solanum. torvum* cultures. Kinetin (Kn) in the concentration of  $2\text{-}3\text{mg l}^{-1}$  induced direct development of shoots from nodal and shoot-tip segments in *Solanum torvum*, optimum response was obtained on  $2\text{mg l}^{-1}$  Kn. The frequency of shoot regeneration was better in nodal culture of *Solanum torvum* than shoot-tip culture. No callus formation was obtained on Kn supported media in nodal and shoot-tip explants. Kn above  $3\text{mg l}^{-1}$  had adverse effect on shoot regeneration in nodal and shoot-tip cultures of *Solanum torvum* [14,15].


In the present investigation, the best shoot regeneration in nodal explant was obtained on  $4\text{mg l}^{-1}$  Kn +  $2\text{mg l}^{-1}$  NAA and in shoot-tip explants on  $3\text{mg l}^{-1}$  Kn +  $2\text{mg l}^{-1}$  NAA and  $2\text{mg l}^{-1}$  2,4-D +  $2\text{mg l}^{-1}$  Kn. Shoot regeneration through callus subculture was frequent in the present experimental system on NAA/ 2,4-D and Kn supplemented media, the optimum response with better shoot regeneration from callus was noted at on  $5\text{mg l}^{-1}$  Kn and  $2\text{mg l}^{-1}$  NAA and  $2\text{mg l}^{-1}$  2,4-D +  $4\text{mg l}^{-1}$  Kn. This is also in conformity of the above facts. Thus, a fine balance of exogenous auxin and cytokinin / cytokinin alone is necessary before successful regeneration can occur.

Kn in combination with NAA / 2,4-D proved effective for shoot regeneration and callus growth in this experimental system [16,17]. The callus in general was greenish-white / white, compact, hydrated and crystalline in appearance. However, in some hormonal combinations, the node derived callus was creamy, white, compact, hydrated and crystalline in appearance. Callus mediated regeneration was frequent in sub culture on  $2\text{mg l}^{-1}$  NAA and  $3\text{-}5\text{mg l}^{-1}$  Kn / 2,4-D + Kn. The optimum response of callus mediated callogenesis was recorded on  $5\text{mg l}^{-1}$  Kn +  $2\text{mg l}^{-1}$  NAA and  $2\text{mg l}^{-1}$  2,4-D +  $4\text{mg l}^{-1}$  Kn. In addition to direct shoot regeneration in nodal and shoot-tip explants, protocol for callus mediated shoot regeneration can also be adopted in the present experimental system as these shoots were morphologically identical to parent plants. The rejuvenation in callus subculture was recorded on  $2\text{mg l}^{-1}$  2,4-D +  $2\text{mg l}^{-1}$  Kn, the callus gradually turned brown in the beginning and after a month profuse shining white callus grew from degenerated callus mass on the same combination of hormones. Calli were maintained at  $25 \pm 2^\circ\text{C}$  in culture till  $1\frac{1}{2}$  years for regeneration on  $1\text{mg l}^{-1}$  NAA +  $1\text{mg l}^{-1}$  Kn and no regeneration was noted on maintenance medium ( $1\text{mg l}^{-1}$  NAA +  $1\text{mg l}^{-1}$  Kn). The rejuvenation in callus subculture was recorded on  $2\text{mg l}^{-1}$  2,4-D +  $2\text{mg l}^{-1}$  Kn, the callus gradually turned brown in the beginning and after a month profuse shining white callus grew from degenerated callus mass on the same combination of hormones.

The best response for shoot regeneration was obtained at  $25 \pm 2^\circ\text{C}$  on  $2\text{mg l}^{-1}$  Kn in nodal and shoot-tip cultures of *Solanum torvum* whereas NAA ( $2\text{mg l}^{-1}$ ) + Kn ( $2\text{-}4\text{mg l}^{-1}$ ) and 2,4-D ( $1\text{-}2\text{mg l}^{-1}$ ) + Kn ( $1\text{-}2\text{mg l}^{-1}$ ) were most responsive combinations for callus growth as well as shoot formation in the present system. In general, auxin and cytokinin above  $5\text{mg l}^{-1}$  were found to be inhibitory for differentiation and regeneration [18]. Direct formation of shoots was frequent in primary cultures of nodal and shoot-tip segments in *Solanum torvum*. Differentiation of callus was obtained on 2,4-D and NAA / 2,4-D + Kn combinations from nodal, internodal, leaf and shoot-tip cultures, callus mediated shoot regeneration was frequent in culture during the present investigation. Identical effects of auxin-



**Figure 1:** *Solanum torvum* Swartz plant growing in nature.



**Figure 2:** Explant showing development callus from entire surface on MS+2mg/l<sup>-1</sup> 2,4-D+2mg/l<sup>-1</sup>Kn; 18 days old culture..



**Figure 3:** 25 days old culture on MS+2mg/l<sup>-1</sup> NAA+3mg/l<sup>-1</sup>Kn.



**Figure 4:** Culture showing development of callus mediated green shoots on MS+2mg/l<sup>-1</sup>NAA+5mg/l<sup>-1</sup>Kn; 25 days old culture.



**Figure 5:** Culture showing development of profuse rooting; 25 days old culture.



**Figure 6:** *In vitro* plant transferred in garden soil.

cytokinin combination on shoot multiplication / callus induction were observed in many plants [19,20]. The role of sugars in tissue culture experiments was extensively studied. They provide energy source and maintain a minimum osmotic potential for the cultured tissue. Many carbohydrates have been used in tissue culture but among them sucrose is the most effective except in a few plants where glucose was found superior to sucrose [21]. Sucrose is generally used at the concentrations of 20-30 g/l and in the present investigation, 30 g/l sucrose was found most effective.

### Conclusion

It was noticed that particular combination of medium was effective for regeneration and growth of callus. Among phytohormones, cytokinin was found to be more pronounced than auxin for callus formation and callus was crystalline. The nodal culture was found to be better than shoot tip culture.

### Acknowledgements

The authors are extremely grateful to Dr. Santosh Kumar, Project Coordinator, UGC-SAP (DRS-Phase-I), University Department of Botany, B.R.A. Bihar University, Muzaffarpur for providing all facilities. The authors feel highly indebted to Dr. Arvind Kumar Jha whose comments and suggestions substantially improved the original manuscript.

### References

1. Bhojwani SS, Razdan, MK (2004) Haploid production In: Plant Tissue Culture-Theory and Practice. (The Amsterdam, Netherlands: Elsevier Science Publisher B.V) pp. 167-213.



2. Garcia-Gonzales R, Quiroz K, Carrasco B, Caligari P (2010) Plant tissue culture: Current status, opportunities and challenges. *Cienc Inv Agr* 37: 5-30.
3. Jaiswal BS (2012) *Solanum torvum*: A review of its traditional uses, phytochemistry and pharmacology. *Int J Pharm Bio Sci* 3: 104-111.
4. CPR Environmental Education Centre (2001) Medicinal plants.
5. Haque A, Islam R, Arima S (2000) High frequency plant regeneration from cotyledon derived callus of *Momordica dioica* (Roxb) wild. *Phytomorphology* 50: 267-272.
6. Rao IU (2008) Understanding in vitro flowering of bamboos by using plant growth regulators; Proceedings of Golden Jubilee Conference on "Challenging and Emerging Strategies for Improving Plant Productivity" (ISPP). IARI, New Delhi, India pp. 52-53.
7. Murashige T, Skoog F (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Plant Physiol* 15: 473-497.
8. Badola KC, Mohinder P, Bhandari HCS, Pal M (1993) Vegetative propagation of banbaigan (*Solanum torvum* Sw.) by rooting branch cuttings. *Indian Forester* 199: 1027-1028.
9. Behera PR, Nayak P, Thirunavoukkarasu M, Sahoo S (2008) Plant regeneration of *Gmelina arborea* Roxb. From cotyledonary node explants. *Indian J Plant Physiol* 13: 258-265.
10. Jagannathan V (1988) Present status of plant tissue culture in India; Proceedings of the National Seminar on Plant Tissue Culture. pp 1-10 (New Delhi: ICAR Publication).
11. Boxus P (1974) The production of strawberry plants by in vitro micropropagation. *J. Hort, Sci* 49: 209-210.
12. Mathew D, Prasad MC (2007) Multiple shoot and plant regeneration from immature leaflets of In vitro origin in curry leaf (*Murraya koenigii*). *Indian J. Plant Physiol* 12: 18-22.
13. Kumari MV, Shivanna MB (2005) Callus mediated regeneration of *Desmodium oojeinensis* Roxb. *Phytomorphology* 55: 166-171.
14. Shanthi P, Annexavier SR (2006) Micropropagation of *Acmellacalva* (DC) R.K. Jansen from nodal explants. *Indian J. Plant Physiol* 11: 89-99.
15. Rathore JS, Rathore MS, Shekhawat NS (2005) Micropropagation of *Maerua oblongifolia*- A liana of arid areas. *Phytomorphology* 55: 241-247.
16. Nair LG, Seeni S (2002) Rapid clonal multiplication of *Morinda umbellata* Linn. (Rubiaceae), A medicinal liana, through cultures of nodes and shoot tips from mature plant. *Phytomorphology* 52: 77-81.
17. Sivasubramanian S, Vallinayagam S, Patric RD, Manicham VS (2002) Micropropagation of *Plectranthus vetiveroides* (Jacob) - A medicinal plant. *Phytomorphology* 52: 55-59.
18. Shahzad A, Siddiqui SA (2000) In vitro organogenesis in *Ocimum sanctum* L.; A multipurpose herb. *Phytomorphology* 50: 27-35.
19. Sharma A, Patna V, Arora DK, Kant U (2008) Micropropagation of *Adenium obesum* L. through apical bud explants - A caudiciform succulent ornamental plant. *J Indian Bot Soc* 87: 232-236.
20. Verma KC, Gaur AK, Singh US (2008) Evaluation of in vitro response from different explant of elite *Jatropha curcas* L. *Indian J Plant Physiol* 13: 231-237.
21. Sreelatha VR, Prasad PJN, Karuppusamy S, Pullaiah T (2008) Morphogenic response of callus derived from hypocotyl explant of *Cassia siamea*. *J Indian Bot Soc* 87: 92-98.